

CS675: Convex and Combinatorial Optimization
Fall 2026
Introduction to Linear Programming

Instructor: Shaddin Dughmi

Outline

- 1 Linear Programming Basics
- 2 Duality and Its Interpretations
- 3 Properties of Duals
- 4 Weak and Strong Duality
- 5 Consequences of Duality
- 6 Formal Proof of Strong Duality of LP
- 7 More Examples of Duality

Outline

- 1 Linear Programming Basics
- 2 Duality and Its Interpretations
- 3 Properties of Duals
- 4 Weak and Strong Duality
- 5 Consequences of Duality
- 6 Formal Proof of Strong Duality of LP
- 7 More Examples of Duality

A Brief History

- The forefather of convex optimization problems, and the most ubiquitous.
- Developed by Kantorovich during World War II (1939) for planning the Soviet army's expenditures and returns. Kept secret.
- Discovered a few years later by George Dantzig, who in 1947 developed the simplex method for solving linear programs
- John von Neumann developed LP duality in 1947, and applied it to game theory
- Polynomial-time algorithms: Ellipsoid method (Khachiyan 1979), interior point methods (Karmarkar 1984).

LP General Form

$$\begin{array}{ll}\text{minimize (or maximize)} & \langle c, x \rangle \\ \text{subject to} & \langle a_i, x \rangle \leq b_i, \quad \text{for } i \in \mathcal{C}^1. \\ & \langle a_i, x \rangle \geq b_i, \quad \text{for } i \in \mathcal{C}^2. \\ & \langle a_i, x \rangle = b_i, \quad \text{for } i \in \mathcal{C}^3.\end{array}$$

- Decision variables: $x \in \mathbb{R}^n$
- Parameters:
 - $c \in \mathbb{R}^n$ defines the linear objective function
 - $a_i \in \mathbb{R}^n$ and $b_i \in \mathbb{R}$ define the i 'th constraint.

LP General Form

$$\begin{array}{ll}\text{minimize (or maximize)} & \langle c, x \rangle \\ \text{subject to} & \langle a_i, x \rangle \leq b_i, \quad \text{for } i \in \mathcal{C}^1. \\ & \langle a_i, x \rangle \geq b_i, \quad \text{for } i \in \mathcal{C}^2. \\ & \langle a_i, x \rangle = b_i, \quad \text{for } i \in \mathcal{C}^3.\end{array}$$

- Decision variables: $x \in \mathbb{R}^n$
- Parameters:
 - $c \in \mathbb{R}^n$ defines the linear objective function
 - $a_i \in \mathbb{R}^n$ and $b_i \in \mathbb{R}$ define the i 'th constraint.

Note

For the inner product (a.k.a. dot product) of vectors u and v I often write $\langle u, v \rangle$, but we can also write $u^\top v$ or $u \cdot v$. Whatever you prefer, or looks most elegant to you.


Maximization Standard Form

$$\begin{array}{ll}\text{maximize} & \langle c, x \rangle \\ \text{subject to} & \langle a_i, x \rangle \leq b_i, \quad \text{for } i = 1, \dots, m. \\ & x_j \geq 0, \quad \text{for } j = 1, \dots, n.\end{array}$$


Every LP can be transformed to this form

- minimizing $\langle c, x \rangle$ is equivalent to maximizing $-\langle c, x \rangle$
- $\geq$ constraints can be flipped by multiplying by -1
- Each equality constraint can be replaced by two inequalities
- Unconstrained variable x_j can be replaced by $x_j^+ - x_j^-$, where both x_j^+ and x_j^- are constrained to be nonnegative.

Geometric View




Geometric View



A 2-D example

$$\begin{array}{ll}\text{maximize} & x_1 + x_2 \\ \text{subject to} & x_1 + 2x_2 \leq 2 \\ & 2x_1 + x_2 \leq 2 \\ & x_1, x_2 \geq 0\end{array}$$



Application: Optimal Production

- n products, m raw materials
- Every unit of product j uses a_{ij} units of raw material i
- There are b_i units of material i available
- Product j yields profit c_j per unit
- Facility wants to maximize profit subject to available raw materials

$$\begin{array}{ll}\text{maximize} & \langle c, x \rangle \\ \text{subject to} & \langle a_i, x \rangle \leq b_i, \quad \text{for } i = 1, \dots, m. \\ & x_j \geq 0, \quad \text{for } j = 1, \dots, n.\end{array}$$


Minimization Standard Form

$$\begin{array}{ll}\text{minimize} & \langle c, x \rangle \\ \text{subject to} & \langle a_i, x \rangle \geq b_i, \quad \text{for } i = 1, \dots, m. \\ & x_j \geq 0, \quad \text{for } j = 1, \dots, n.\end{array}$$

Every LP can be transformed to this form similarly

Terminology

- **Hyperplane**: The region defined by a linear equality
- **Halfspace**: The region defined by a linear inequality $\langle a_i, x \rangle \leq b_i$.
- **Polyhedron**: The intersection of a set of linear inequalities
 - Feasible region of an LP is a polyhedron
- **Polytope**: Bounded polyhedron
 - Equivalently: **convex hull** of a finite set of points
- **Vertex**: A point x is a vertex of polyhedron P if $\nexists y \neq 0$ with $x + y \in P$ and $x - y \in P$
- **Face** of P : The intersection with P of a hyperplane H disjoint from the interior of P



Basic Facts about LPs and Polyhedrons

Fact

Feasible regions of LPs (i.e. polyhedrons) are convex

Basic Facts about LPs and Polyhedrons

Fact

Feasible regions of LPs (i.e. polyhedrons) are convex

Fact

Set of optimal solutions of an LP is convex

- In fact, a face of the polyhedron
- intersection of P with hyperplane $\langle c, x \rangle = OPT$

Basic Facts about LPs and Polyhedrons

Fact

Feasible regions of LPs (i.e. polyhedrons) are convex

Fact

Set of optimal solutions of an LP is convex

- In fact, a face of the polyhedron
- intersection of P with hyperplane $\langle c, x \rangle = OPT$



Fact

A feasible point x is a vertex if and only if n linearly independent constraints are **tight** (i.e., satisfied with equality) at x .

Basic Facts about LPs and Polyhedrons

Fact

An LP either has an optimal solution, or is **unbounded** or **infeasible**



Fundamental Theorem of LP

If an LP in standard form has an optimal solution, then it has a vertex optimal solution.

Fundamental Theorem of LP

If an LP in standard form has an optimal solution, then it has a vertex optimal solution.

Proof

- Assume not, and take a non-vertex optimal solution x with the maximum number of tight constraints

Fundamental Theorem of LP

If an LP in standard form has an optimal solution, then it has a vertex optimal solution.

Proof

- Assume not, and take a non-vertex optimal solution x with the maximum number of tight constraints
- There is $y \neq 0$ s.t. $x \pm y$ are feasible

Fundamental Theorem of LP

If an LP in standard form has an optimal solution, then it has a vertex optimal solution.

Proof

- Assume not, and take a non-vertex optimal solution x with the maximum number of tight constraints
- There is $y \neq 0$ s.t. $x \pm y$ are feasible
- y is perpendicular to the objective function and the tight constraints at x .
 - i.e. $\langle c, y \rangle = 0$, and $\langle a_i, y \rangle = 0$ whenever i 'th constraint is tight for x .

Fundamental Theorem of LP

If an LP in standard form has an optimal solution, then it has a vertex optimal solution.

Proof

- Assume not, and take a non-vertex optimal solution x with the maximum number of tight constraints
- There is $y \neq 0$ s.t. $x \pm y$ are feasible
- y is perpendicular to the objective function and the tight constraints at x .
 - i.e. $\langle c, y \rangle = 0$, and $\langle a_i, y \rangle = 0$ whenever i 'th constraint is tight for x .
- Can choose y s.t. $y_j < 0$ for some j

Fundamental Theorem of LP

If an LP in standard form has an optimal solution, then it has a vertex optimal solution.

Proof

- Assume not, and take a non-vertex optimal solution x with the maximum number of tight constraints
- There is $y \neq 0$ s.t. $x \pm y$ are feasible
- y is perpendicular to the objective function and the tight constraints at x .
 - i.e. $\langle c, y \rangle = 0$, and $\langle a_i, y \rangle = 0$ whenever i 'th constraint is tight for x .
- Can choose y s.t. $y_j < 0$ for some j
- Let α be the largest constant such that $x + \alpha y$ is feasible
 - Such an α exists

Fundamental Theorem of LP

If an LP in standard form has an optimal solution, then it has a vertex optimal solution.

Proof

- Assume not, and take a non-vertex optimal solution x with the maximum number of tight constraints
- There is $y \neq 0$ s.t. $x \pm y$ are feasible
- y is perpendicular to the objective function and the tight constraints at x .
 - i.e. $\langle c, y \rangle = 0$, and $\langle a_i, y \rangle = 0$ whenever i 'th constraint is tight for x .
- Can choose y s.t. $y_j < 0$ for some j
- Let α be the largest constant such that $x + \alpha y$ is feasible
 - Such an α exists
- An additional constraint becomes tight at $x + \alpha y$, a contradiction.

Counting non-zero Variables

Corollary

If an LP in standard form has an optimal solution, then there is an optimal solution with at most m non-zero variables.

$$\begin{array}{ll}\text{maximize} & \langle c, x \rangle \\ \text{subject to} & \langle a_i, x \rangle \leq b_i, \quad \text{for } i = 1, \dots, m. \\ & x_j \geq 0, \quad \text{for } j = 1, \dots, n.\end{array}$$

- e.g. for optimal production with n products and m raw materials, there is an optimal plan with at most m products.

Fundamental Theorem of LP (General Version)

If an LP has an optimal solution, and moreover it's feasible region includes no lines, then it has a vertex optimal solution.

Fundamental Theorem of LP (General Version)

If an LP has an optimal solution, and moreover it's feasible region includes no lines, then it has a vertex optimal solution.

- Essentially the same proof (exercise)
- In addition to LPs in standard form, applies to LPs with a bounded feasible region (i.e., feasible region is a polytope), among others.

Outline

- 1 Linear Programming Basics
- 2 Duality and Its Interpretations**
- 3 Properties of Duals
- 4 Weak and Strong Duality
- 5 Consequences of Duality
- 6 Formal Proof of Strong Duality of LP
- 7 More Examples of Duality

Linear Programming Duality

Primal LP

$$\begin{array}{ll}\text{maximize} & \langle c, x \rangle \\ \text{subject to} & Ax \preceq b\end{array}$$

Dual LP

$$\begin{array}{ll}\text{minimize} & \langle b, y \rangle \\ \text{subject to} & A^T y = c \\ & y \succeq 0\end{array}$$

- $A \in \mathbb{R}^{m \times n}$, $c \in \mathbb{R}^n$, $b \in \mathbb{R}^m$
- y_i is the **dual variable** corresponding to primal constraint $A_i x \leq b_i$
- $A_j^T y = c_j$ is the **dual constraint** corresponding to primal variable x_j

Linear Programming Duality: Standard Form, and Visualization

Primal LP

maximize $\langle c, x \rangle$
subject to $Ax \preceq b$
 $x \succeq 0$

Dual LP

minimize $\langle b, y \rangle$
subject to $A^T y \succeq c$
 $y \succeq 0$

Linear Programming Duality: Standard Form, and Visualization

Primal LP

maximize $\langle c, x \rangle$
subject to $Ax \preceq b$
 $x \succeq 0$

Dual LP

minimize $\langle b, y \rangle$
subject to $A^T y \succeq c$
 $y \succeq 0$

	x_1	x_2	x_3	x_4	
y_1	a_{11}	a_{12}	a_{13}	a_{14}	b_1
y_2	a_{21}	a_{22}	a_{23}	a_{24}	b_2
y_3	a_{31}	a_{32}	a_{33}	a_{34}	b_3
	c_1	c_2	c_3	c_4	

Linear Programming Duality: Standard Form, and Visualization

Primal LP

$$\begin{array}{ll}\text{maximize} & \langle c, x \rangle \\ \text{subject to} & Ax \preceq b \\ & x \succeq 0\end{array}$$

Dual LP

$$\begin{array}{ll}\text{minimize} & \langle b, y \rangle \\ \text{subject to} & A^T y \succeq c \\ & y \succeq 0\end{array}$$

	x_1	x_2	x_3	x_4	
y_1	a_{11}	a_{12}	a_{13}	a_{14}	b_1
y_2	a_{21}	a_{22}	a_{23}	a_{24}	b_2
y_3	a_{31}	a_{32}	a_{33}	a_{34}	b_3
	c_1	c_2	c_3	c_4	

- y_i is the **dual variable** corresponding to primal constraint $A_i x \leq b_i$
- $A_j^T y \geq c_j$ is the **dual constraint** corresponding to primal variable x_j

Interpretation 1: Economic Interpretation

Recall the Optimal Production problem from last lecture

- n products, m raw materials
- Every unit of product j uses a_{ij} units of raw material i
- There are b_i units of material i available
- Product j yields profit c_j per unit
- Facility wants to maximize profit subject to available raw materials

Interpretation 1: Economic Interpretation

Primal LP

$$\begin{array}{ll}\max & \sum_{j=1}^n c_j x_j \\ \text{s.t.} & \sum_{j=1}^n a_{ij} x_j \leq b_i, \quad \text{for } i \in [m]. \\ & x_j \geq 0, \quad \text{for } j \in [n].\end{array}$$

Interpretation 1: Economic Interpretation

Primal LP

$$\begin{array}{ll}\max & \sum_{j=1}^n c_j x_j \\ \text{s.t.} & \sum_{j=1}^n a_{ij} x_j \leq b_i, \quad \text{for } i \in [m]. \\ & x_j \geq 0, \quad \text{for } j \in [n].\end{array}$$

Dual LP

$$\begin{array}{ll}\min & \sum_{i=1}^m b_i y_i \\ \text{s.t.} & \sum_{i=1}^m a_{ij} y_i \geq c_j, \quad \text{for } j \in [n]. \\ & y_i \geq 0, \quad \text{for } i \in [m].\end{array}$$

Interpretation 1: Economic Interpretation

Primal LP

$$\begin{array}{ll}\max & \sum_{j=1}^n c_j x_j \\ \text{s.t.} & \sum_{j=1}^n a_{ij} x_j \leq b_i, \quad \text{for } i \in [m]. \\ & x_j \geq 0, \quad \text{for } j \in [n].\end{array}$$

Dual LP

$$\begin{array}{ll}\min & \sum_{i=1}^m b_i y_i \\ \text{s.t.} & \sum_{i=1}^m a_{ij} y_i \geq c_j, \quad \text{for } j \in [n]. \\ & y_i \geq 0, \quad \text{for } i \in [m].\end{array}$$

	x_1	x_2	x_3	x_4	
y_1	a_{11}	a_{12}	a_{13}	a_{14}	b_1
y_2	a_{21}	a_{22}	a_{23}	a_{24}	b_2
y_3	a_{31}	a_{32}	a_{33}	a_{34}	b_3
	c_1	c_2	c_3	c_4	

Interpretation 1: Economic Interpretation

Primal LP

$$\begin{array}{ll}\max & \sum_{j=1}^n c_j x_j \\ \text{s.t.} & \sum_{j=1}^n a_{ij} x_j \leq b_i, \quad \text{for } i \in [m]. \\ & x_j \geq 0, \quad \text{for } j \in [n].\end{array}$$

Dual LP

$$\begin{array}{ll}\min & \sum_{i=1}^m b_i y_i \\ \text{s.t.} & \sum_{i=1}^m a_{ij} y_i \geq c_j, \quad \text{for } j \in [n]. \\ & y_i \geq 0, \quad \text{for } i \in [m].\end{array}$$

	x_1	x_2	x_3	x_4	
y_1	a_{11}	a_{12}	a_{13}	a_{14}	b_1
y_2	a_{21}	a_{22}	a_{23}	a_{24}	b_2
y_3	a_{31}	a_{32}	a_{33}	a_{34}	b_3
	c_1	c_2	c_3	c_4	

- Dual variable y_i is a proposed **price** per unit of raw material i
- Dual price vector is feasible if facility has incentive to sell materials
- Buyer wants to spend as little as possible to buy materials

Interpretation 2: Finding the Best Upperbound

Consider the simple LP from last lecture

$$\begin{array}{ll}\text{maximize} & x_1 + x_2 \\ \text{subject to} & x_1 + 2x_2 \leq 2 \\ & 2x_1 + x_2 \leq 2 \\ & x_1, x_2 \geq 0\end{array}$$

- We found that the optimal solution was at $(\frac{2}{3}, \frac{2}{3})$, with an optimal value of $4/3$.

Interpretation 2: Finding the Best Upperbound

Consider the simple LP from last lecture

$$\begin{array}{ll}\text{maximize} & x_1 + x_2 \\ \text{subject to} & x_1 + 2x_2 \leq 2 \\ & 2x_1 + x_2 \leq 2 \\ & x_1, x_2 \geq 0\end{array}$$

- We found that the optimal solution was at $(\frac{2}{3}, \frac{2}{3})$, with an optimal value of $4/3$.
- What if, instead of finding the optimal solution, we sought to find an upperbound on its value by combining inequalities?
 - Each inequality implies an upper bound of 2
 - Multiplying each by $\frac{1}{3}$ and summing gives $x_1 + x_2 \leq 4/3$.

Interpretation 2: Finding the Best Upperbound

	x_1	x_2	x_3	x_4	
y_1	a_{11}	a_{12}	a_{13}	a_{14}	b_1
y_2	a_{21}	a_{22}	a_{23}	a_{24}	b_2
y_3	a_{31}	a_{32}	a_{33}	a_{34}	b_3
	c_1	c_2	c_3	c_4	

- Multiplying each row i by y_i and summing gives the inequality

$$y^T A x \leq y^T b$$

Interpretation 2: Finding the Best Upperbound

	x_1	x_2	x_3	x_4	
y_1	a_{11}	a_{12}	a_{13}	a_{14}	b_1
y_2	a_{21}	a_{22}	a_{23}	a_{24}	b_2
y_3	a_{31}	a_{32}	a_{33}	a_{34}	b_3
	c_1	c_2	c_3	c_4	

- Multiplying each row i by y_i and summing gives the inequality

$$y^T A x \leq y^T b$$

- When $y^T A \geq c^T$, the right hand side of the inequality is an upper bound on $c^T x$ for every feasible x .

$$c^T x \leq y^T A x \leq y^T b$$

Interpretation 2: Finding the Best Upperbound

	x_1	x_2	x_3	x_4	
y_1	a_{11}	a_{12}	a_{13}	a_{14}	b_1
y_2	a_{21}	a_{22}	a_{23}	a_{24}	b_2
y_3	a_{31}	a_{32}	a_{33}	a_{34}	b_3
	c_1	c_2	c_3	c_4	

- Multiplying each row i by y_i and summing gives the inequality


$$y^T A x \leq y^T b$$

- When $y^T A \geq c^T$, the right hand side of the inequality is an upper bound on $c^T x$ for every feasible x .

$$c^T x \leq y^T A x \leq y^T b$$


- The dual LP can be thought of as trying to find the best upperbound on the primal that can be achieved this way.

Interpretation 3: Physical Forces




- Apply force field c to a ball inside bounded polytope $Ax \preceq b$.

Interpretation 3: Physical Forces




- Apply force field c to a ball inside bounded polytope $Ax \preceq b$.
- Eventually, ball will come to rest against the walls of the polytope.

Interpretation 3: Physical Forces




- Apply force field c to a ball inside bounded polytope $Ax \preceq b$.
- Eventually, ball will come to rest against the walls of the polytope.
- Wall $a_i x \leq b_i$ applies some force $-y_i a_i$ to the ball

Interpretation 3: Physical Forces



- Apply force field c to a ball inside bounded polytope $Ax \preceq b$.
- Eventually, ball will come to rest against the walls of the polytope.
- Wall $a_i x \leq b_i$ applies some force $-y_i a_i$ to the ball
- Since the ball is still, $c^T = \sum_i y_i a_i = y^T A$.

Interpretation 3: Physical Forces



- Apply force field c to a ball inside bounded polytope $Ax \preceq b$.
- Eventually, ball will come to rest against the walls of the polytope.
- Wall $a_i x \leq b_i$ applies some force $-y_i a_i$ to the ball
- Since the ball is still, $c^T = \sum_i y_i a_i = y^T A$.
- Dual can be thought of as trying to minimize “work” $\sum_i y_i b_i$ to bring ball back to origin by moving polytope
- We will see that, at optimality, only the walls adjacent to the ball push (Complementary Slackness)

Outline

- 1 Linear Programming Basics
- 2 Duality and Its Interpretations
- 3 Properties of Duals**
- 4 Weak and Strong Duality
- 5 Consequences of Duality
- 6 Formal Proof of Strong Duality of LP
- 7 More Examples of Duality

Duality is an Inversion

Primal LP

$$\begin{array}{ll}\text{maximize} & \langle c, x \rangle \\ \text{subject to} & Ax \preceq b \\ & x \succeq 0\end{array}$$

Dual LP

$$\begin{array}{ll}\text{minimize} & \langle b, y \rangle \\ \text{subject to} & A^T y \succeq c \\ & y \succeq 0\end{array}$$

Duality is an Inversion

Given a primal LP in standard form, the dual of its dual is itself.

Correspondance Between Variables and Constraints

Primal LP

$$\begin{array}{ll}\max & \sum_{j=1}^n c_j x_j \\ \text{s.t.} & \sum_{j=1}^n a_{ij} x_j \leq b_i, \quad \text{for } i \in [m]. \\ & x_j \geq 0, \quad \text{for } j \in [n].\end{array}$$

Dual LP

$$\begin{array}{ll}\min & \sum_{i=1}^m b_i y_i \\ \text{s.t.} & \sum_{i=1}^m a_{ij} y_i \geq c_j, \quad \text{for } j \in [n]. \\ & y_i \geq 0, \quad \text{for } i \in [m].\end{array}$$

Correspondance Between Variables and Constraints

Primal LP

$$\begin{array}{ll}\max & \sum_{j=1}^n c_j x_j \\ \text{s.t.} & \\ \textcolor{red}{y_i} : & \sum_{j=1}^n a_{ij} x_j \leq b_i, \quad \text{for } i \in [m]. \\ & x_j \geq 0, \quad \text{for } j \in [n].\end{array}$$

Dual LP

$$\begin{array}{ll}\min & \sum_{i=1}^m b_i y_i \\ \text{s.t.} & \\ & \sum_{i=1}^m a_{ij} y_i \geq c_j, \quad \text{for } j \in [n]. \\ & y_i \geq 0, \quad \text{for } i \in [m].\end{array}$$

- The i 'th primal constraint gives rise to the i 'th dual variable y_i

Correspondance Between Variables and Constraints

Primal LP

$$\begin{array}{ll}\max & \sum_{j=1}^n c_j x_j \\ \text{s.t.} & \\ \textcolor{red}{y_i} : & \sum_{j=1}^n a_{ij} x_j \leq b_i, \quad \text{for } i \in [m]. \\ & x_j \geq 0, \quad \text{for } j \in [n].\end{array}$$

Dual LP

$$\begin{array}{ll}\min & \sum_{i=1}^m b_i y_i \\ \text{s.t.} & \\ \textcolor{red}{x_j} : & \sum_{i=1}^m a_{ij} y_i \geq c_j, \quad \text{for } j \in [n]. \\ & y_i \geq 0, \quad \text{for } i \in [m].\end{array}$$

- The i 'th primal constraint gives rise to the i 'th dual variable y_i
- The j 'th primal variable x_j gives rise to the j 'th dual constraint

Syntactic Rules

Primal LP

$$\begin{array}{ll}\max & \langle c, x \rangle \\ \text{s.t.} & \\ y_i : & a_i x \leq b_i, \quad \text{for } i \in \mathcal{C}_1. \\ y_i : & a_i x = b_i, \quad \text{for } i \in \mathcal{C}_2. \\ & x_j \geq 0, \quad \text{for } j \in \mathcal{D}_1. \\ & x_j \in \mathbb{R}, \quad \text{for } j \in \mathcal{D}_2.\end{array}$$

Dual LP

$$\begin{array}{ll}\min & \langle b, y \rangle \\ \text{s.t.} & \\ x_j : & \langle \bar{a}_j, y \rangle \geq c_j, \quad \text{for } j \in \mathcal{D}_1. \\ x_j : & \langle \bar{a}_j, y \rangle = c_j, \quad \text{for } j \in \mathcal{D}_2. \\ & y_i \geq 0, \quad \text{for } i \in \mathcal{C}_1. \\ & y_i \in \mathbb{R}, \quad \text{for } i \in \mathcal{C}_2.\end{array}$$

Rules of Thumb

- Lenient constraint (i.e. inequality) $\Rightarrow$ stringent dual variable (i.e. nonnegative)
- Stringent constraint (i.e. equality) $\Rightarrow$ lenient dual variable (i.e. unconstrained)

Outline

- 1 Linear Programming Basics
- 2 Duality and Its Interpretations
- 3 Properties of Duals
- 4 Weak and Strong Duality**
- 5 Consequences of Duality
- 6 Formal Proof of Strong Duality of LP
- 7 More Examples of Duality

Weak Duality

Primal LP

maximize $\langle c, x \rangle$
subject to $Ax \preceq b$
 $x \succeq 0$

Dual LP

minimize $\langle b, y \rangle$
subject to $A^T y \succeq c$
 $y \succeq 0$

Theorem (Weak Duality)

For every primal feasible x and dual feasible y , we have $\langle c, x \rangle \leq \langle b, y \rangle$.

Corollary

- *If primal and dual both feasible and bounded, $OPT(Primal) \leq OPT(Dual)$*
- *If primal is unbounded, dual is infeasible*
- *If dual is unbounded, primal is infeasible*

Weak Duality

Primal LP

maximize $\langle c, x \rangle$
subject to $Ax \preceq b$
 $x \succeq 0$

Dual LP

minimize $\langle b, y \rangle$
subject to $A^T y \succeq c$
 $y \succeq 0$

Theorem (Weak Duality)

For every primal feasible x and dual feasible y , we have $\langle c, x \rangle \leq \langle b, y \rangle$.

Corollary

If x^ is primal feasible, and y^* is dual feasible, and $\langle c, x^* \rangle = \langle b, y^* \rangle$, then both are optimal.*

Interpretation of Weak Duality

Economic Interpretation

For prices are feasible for the dual, the total money collected from sale of raw materials would exceed profit from production.

Interpretation of Weak Duality

Economic Interpretation

For prices are feasible for the dual, the total money collected from sale of raw materials would exceed profit from production.

Upperbound Interpretation

The method of scaling and summing inequalities yields a sound proof system for upperbounds on the primal optimal value.

Interpretation of Weak Duality

Economic Interpretation

For prices are feasible for the dual, the total money collected from sale of raw materials would exceed profit from production.

Upperbound Interpretation

The method of scaling and summing inequalities yields a sound proof system for upperbounds on the primal optimal value.

Physical Interpretation

Work required to bring ball back to origin by pulling polytope is at least potential energy difference between origin and primal optimum.

Proof of Weak Duality

Primal LP

maximize $\langle c, x \rangle$
subject to $Ax \preceq b$
 $x \succeq 0$

Dual LP

minimize $\langle b, y \rangle$
subject to $A^T y \succeq c$
 $y \succeq 0$

$$c^T x \leq y^T Ax \leq y^T b$$

Strong Duality

Primal LP

maximize $\langle c, x \rangle$
subject to $Ax \preceq b$
 $x \succeq 0$

Dual LP

minimize $\langle b, y \rangle$
subject to $A^T y \succeq c$
 $y \succeq 0$

Theorem (Strong Duality)

If either the primal or dual is feasible and bounded, then so is the other and $OPT(Primal) = OPT(Dual)$.

Interpretation of Strong Duality

Economic Interpretation

Buyer can offer dual-feasible prices for raw materials that would make facility indifferent between production and sale.

Interpretation of Strong Duality

Economic Interpretation

Buyer can offer dual-feasible prices for raw materials that would make facility indifferent between production and sale.

Upperbound Interpretation

The method of scaling and summing inequalities yields a complete proof system for upperbounds on the primal optimal value.

Interpretation of Strong Duality

Economic Interpretation

Buyer can offer dual-feasible prices for raw materials that would make facility indifferent between production and sale.


Upperbound Interpretation

The method of scaling and summing inequalities yields a complete proof system for upperbounds on the primal optimal value.

Physical Interpretation


There is an assignment of forces to the walls of the polytope that brings ball back to the origin without wasting energy.

Informal Proof of Strong Duality




- Recall the physical interpretation of duality

Informal Proof of Strong Duality



- Recall the physical interpretation of duality
- When ball is stationary at x , we expect force c to be neutralized only by constraints that are tight. i.e. force multipliers $y \succeq 0$ s.t.
 - $y^T A = c$
 - $y_i(b_i - a_i x) = 0$

Informal Proof of Strong Duality



- Recall the physical interpretation of duality
- When ball is stationary at x , we expect force c to be neutralized only by constraints that are tight. i.e. force multipliers $y \succeq 0$ s.t.
 - $y^T A = c$
 - $y_i(b_i - a_i x) = 0$

$$y^T b - c^T x = y^T b - y^T A x = \sum_i y_i(b_i - a_i x) = 0$$

We found a primal and dual solution that are equal in value!

Outline

- 1 Linear Programming Basics
- 2 Duality and Its Interpretations
- 3 Properties of Duals
- 4 Weak and Strong Duality
- 5 Consequences of Duality**
- 6 Formal Proof of Strong Duality of LP
- 7 More Examples of Duality

Complementary Slackness

Primal LP

maximize $\langle c, x \rangle$
subject to $Ax \preceq b$
 $x \succeq 0$

Dual LP

minimize $\langle y, b \rangle$
subject to $A^T y \succeq c$
 $y \succeq 0$

Complementary Slackness

Primal LP

$$\begin{array}{ll}\text{maximize} & \langle c, x \rangle \\ \text{subject to} & Ax \preceq b \\ & x \succeq 0\end{array}$$

Dual LP

$$\begin{array}{ll}\text{minimize} & \langle y, b \rangle \\ \text{subject to} & A^T y \succeq c \\ & y \succeq 0\end{array}$$

- Let $s_i = (b - Ax)_i$ be the i 'th **primal slack variable**
- Let $t_j = (A^T y - c)_j$ be the j 'th **dual slack variable**

Complementary Slackness

Primal LP

$$\begin{array}{ll}\text{maximize} & \langle c, x \rangle \\ \text{subject to} & Ax \preceq b \\ & x \succeq 0\end{array}$$

Dual LP

$$\begin{array}{ll}\text{minimize} & \langle y, b \rangle \\ \text{subject to} & A^T y \succeq c \\ & y \succeq 0\end{array}$$

- Let $s_i = (b - Ax)_i$ be the i 'th **primal slack variable**
- Let $t_j = (A^T y - c)_j$ be the j 'th **dual slack variable**

Complementary Slackness

Feasible x and y are optimal if and only if

- $x_j t_j = 0$ for all $j = 1, \dots, n$
- $y_i s_i = 0$ for all $i = 1, \dots, m$

	x_1	x_2	x_3	x_4	
y_1	a_{11}	a_{12}	a_{13}	a_{14}	b_1
y_2	a_{21}	a_{22}	a_{23}	a_{24}	b_2
y_3	a_{31}	a_{32}	a_{33}	a_{34}	b_3
	c_1	c_2	c_3	c_4	

Interpretation of Complementary Slackness

Economic Interpretation


Given an optimal primal production vector x and optimal dual offer prices y ,

- Facility produces only products for which it is indifferent between sale and production.
- Only raw materials that are binding constraints on production are priced greater than 0

Interpretation of Complementary Slackness

Physical Interpretation

Only walls adjacent to the ball's equilibrium position push back on it.



Proof of Complementary Slackness

Primal LP

maximize $\langle c, x \rangle$
subject to $Ax \preceq b$
 $x \succeq 0$

Dual LP

minimize $\langle y, b \rangle$
subject to $A^T y \succeq c$
 $y \succeq 0$

Proof of Complementary Slackness

Primal LP

$$\begin{array}{ll}\text{maximize} & \langle c, x \rangle \\ \text{subject to} & Ax + s = b \\ & x \succeq 0 \\ & s \succeq 0\end{array}$$

Dual LP

$$\begin{array}{ll}\text{minimize} & \langle y, b \rangle \\ \text{subject to} & A^T y - t = c \\ & y \succeq 0 \\ & t \succeq 0\end{array}$$

- Can equivalently rewrite LP using slack variables

Proof of Complementary Slackness

Primal LP

$$\begin{array}{ll}\text{maximize} & \langle c, x \rangle \\ \text{subject to} & Ax + s = b \\ & x \succeq 0 \\ & s \succeq 0\end{array}$$

Dual LP

$$\begin{array}{ll}\text{minimize} & \langle y, b \rangle \\ \text{subject to} & A^T y - t = c \\ & y \succeq 0 \\ & t \succeq 0\end{array}$$

- Can equivalently rewrite LP using slack variables

$$y^T b - c^T x = y^T (Ax + s) - (y^T A - t^T) x = y^T s + t^T x$$

Proof of Complementary Slackness

Primal LP

$$\begin{array}{ll}\text{maximize} & \langle c, x \rangle \\ \text{subject to} & Ax + s = b \\ & x \succeq 0 \\ & s \succeq 0\end{array}$$

Dual LP

$$\begin{array}{ll}\text{minimize} & \langle y, b \rangle \\ \text{subject to} & A^T y - t = c \\ & y \succeq 0 \\ & t \succeq 0\end{array}$$

- Can equivalently rewrite LP using slack variables

$$y^T b - c^T x = y^T (Ax + s) - (y^T A - t^T)x = y^T s + t^T x$$

Gap between primal and dual objectives is 0 if and only if complementary slackness holds.

Recovering Primal from Dual

- Will encounter LPs where the dual is easier to solve than primal
- Complementary slackness allows us to recover the primal optimal from the dual optimal, and vice versa.

Recovering Primal from Dual

- Will encounter LPs where the dual is easier to solve than primal
- Complementary slackness allows us to recover the primal optimal from the dual optimal, and vice versa.
 - Assuming **non-degeneracy**: At every vertex of primal [dual] there are exactly n [m] tight constraints which are linearly independent.

Recovering Primal from Dual

- Will encounter LPs where the dual is easier to solve than primal
- Complementary slackness allows us to recover the primal optimal from the dual optimal, and vice versa.
 - Assuming **non-degeneracy**: At every vertex of primal [dual] there are exactly n [m] tight constraints which are linearly independent.

Primal LP

(n variables, $m + n$ constraints)

$$\begin{array}{ll}\text{maximize} & \langle c, x \rangle \\ \text{subject to} & Ax \preceq b \\ & x \succeq 0\end{array}$$

Dual LP

(m variables, $m + n$ constraints)

$$\begin{array}{ll}\text{minimize} & \langle y, b \rangle \\ \text{subject to} & A^T y \succeq c \\ & y \succeq 0\end{array}$$

Recovering Primal from Dual

- Will encounter LPs where the dual is easier to solve than primal
- Complementary slackness allows us to recover the primal optimal from the dual optimal, and vice versa.
 - Assuming **non-degeneracy**: At every vertex of primal [dual] there are exactly n [m] tight constraints which are linearly independent.

Primal LP

(n variables, $m + n$ constraints)

$$\begin{array}{ll}\text{maximize} & \langle c, x \rangle \\ \text{subject to} & Ax \preceq b \\ & x \succeq 0\end{array}$$

Dual LP

(m variables, $m + n$ constraints)

$$\begin{array}{ll}\text{minimize} & \langle y, b \rangle \\ \text{subject to} & A^T y \succeq c \\ & y \succeq 0\end{array}$$

- Let y be dual optimal. By non-degeneracy:
 - Exactly m of the $m + n$ dual constraints are tight at y
 - Exactly n dual constraints are loose

Recovering Primal from Dual

- Will encounter LPs where the dual is easier to solve than primal
- Complementary slackness allows us to recover the primal optimal from the dual optimal, and vice versa.
 - Assuming **non-degeneracy**: At every vertex of primal [dual] there are exactly n [m] tight constraints which are linearly independent.

Primal LP

(n variables, $m + n$ constraints)

$$\begin{array}{ll}\text{maximize} & \langle c, x \rangle \\ \text{subject to} & Ax \preceq b \\ & x \succeq 0\end{array}$$

Dual LP

(m variables, $m + n$ constraints)

$$\begin{array}{ll}\text{minimize} & \langle y, b \rangle \\ \text{subject to} & A^T y \succeq c \\ & y \succeq 0\end{array}$$

- Let y be dual optimal. By non-degeneracy:
 - Exactly m of the $m + n$ dual constraints are tight at y
 - Exactly n dual constraints are loose
- n loose dual constraints impose n tight primal constraints

Recovering Primal from Dual

- Will encounter LPs where the dual is easier to solve than primal
- Complementary slackness allows us to recover the primal optimal from the dual optimal, and vice versa.
 - Assuming **non-degeneracy**: At every vertex of primal [dual] there are exactly n [m] tight constraints which are linearly independent.

Primal LP

(n variables, $m + n$ constraints)

$$\begin{array}{ll}\text{maximize} & \langle c, x \rangle \\ \text{subject to} & Ax \preceq b \\ & x \succeq 0\end{array}$$

Dual LP

(m variables, $m + n$ constraints)

$$\begin{array}{ll}\text{minimize} & \langle y, b \rangle \\ \text{subject to} & A^T y \succeq c \\ & y \succeq 0\end{array}$$

- Let y be dual optimal. By non-degeneracy:
 - Exactly m of the $m + n$ dual constraints are tight at y
 - Exactly n dual constraints are loose
- n loose dual constraints impose n tight primal constraints
 - Assuming non-degeneracy, solving the linear equation yields a unique primal optimum solution x .

Sensitivity Analysis

Primal LP

$$\begin{array}{ll}\text{maximize} & \langle c, x \rangle \\ \text{subject to} & Ax \preceq b \\ & x \succeq 0\end{array}$$

Dual LP

$$\begin{array}{ll}\text{minimize} & \langle y, b \rangle \\ \text{subject to} & A^T y \succeq c \\ & y \succeq 0\end{array}$$

Sometimes, we want to examine how the optimal value of our LP changes with its parameters c and b

Sensitivity Analysis

Primal LP

$$\begin{array}{ll}\text{maximize} & \langle c, x \rangle \\ \text{subject to} & Ax \preceq b \\ & x \succeq 0\end{array}$$

Dual LP

$$\begin{array}{ll}\text{minimize} & \langle y, b \rangle \\ \text{subject to} & A^T y \succeq c \\ & y \succeq 0\end{array}$$

Sometimes, we want to examine how the optimal value of our LP changes with its parameters c and b

Sensitivity Analysis

Let $OPT = OPT(A, c, b)$ be the optimal value of the above LP. Let x and y be the primal and dual optima.

- $\frac{\partial OPT}{\partial c_j} = x_j$ when x is the unique primal optimum.
- $\frac{\partial OPT}{\partial b_i} = y_i$ when y is the unique dual optimum.

Sensitivity Analysis

Primal LP

$$\begin{array}{ll}\text{maximize} & \langle c, x \rangle \\ \text{subject to} & Ax \preceq b \\ & x \succeq 0\end{array}$$

Dual LP

$$\begin{array}{ll}\text{minimize} & \langle y, b \rangle \\ \text{subject to} & A^T y \succeq c \\ & y \succeq 0\end{array}$$

Sometimes, we want to examine how the optimal value of our LP changes with its parameters c and b

Economic Interpretation of Sensitivity Analysis


- A small increase δ in c_j increases profit by $\delta \cdot x_j$
- A small increase δ in b_i increases profit by $\delta \cdot y_i$
 - y_i measures the “marginal value” of resource i for production

Outline

- 1 Linear Programming Basics
- 2 Duality and Its Interpretations
- 3 Properties of Duals
- 4 Weak and Strong Duality
- 5 Consequences of Duality
- 6 Formal Proof of Strong Duality of LP**
- 7 More Examples of Duality

Separating Hyperplane Theorem

For any two disjoint convex sets, then there is a hyperplane separating them. Moreover, if both sets are closed and at least one of them is compact, then there is a hyperplane strictly separating them.




Definition

A **convex cone** is a convex subset of $\mathbb{R}^n$ which is closed under nonnegative scaling and convex combinations.

Definition

The convex cone **generated** by vectors $u_1, \dots, u_m \in \mathbb{R}^n$ is the set of all nonnegative-weighted sums of these vectors (also known as **conic combinations**).

$$\text{Cone}(u_1, \dots, u_m) = \left\{ \sum_{i=1}^m \alpha_i u_i : \alpha_i \geq 0 \forall i \right\}$$




This follows from the separating hyperplane thm (exercise).

Farkas' Lemma

Let $\mathcal{C}$ be the **convex cone** generated by vectors $u_1, \dots, u_m \in \mathbb{R}^n$, and let $w \in \mathbb{R}^n$. Exactly one of the following is true:


- $w \in \mathcal{C}$
- There is $z \in \mathbb{R}^n$ such that $z \cdot u_i \leq 0$ for all i , and $z \cdot w > 0$.



Equivalently: Theorem of the Alternative

Exactly one of the following is true for $U = [u_1, \dots, u_m]$ and w

- The system $Uz = w, z \succeq 0$ has a solution
- The system $U^T z \preceq 0, \langle z, w \rangle > 0$ has a solution.



Formal Proof of Strong Duality

Primal LP

$$\begin{array}{ll}\text{maximize} & \langle c, x \rangle \\ \text{subject to} & Ax \preceq b\end{array}$$

Dual LP

$$\begin{array}{ll}\text{minimize} & \langle b, y \rangle \\ \text{subject to} & A^\top y = c \\ & y \succeq 0\end{array}$$

Given $v \in \mathbb{R}$, by Farkas' Lemma exactly one of the following is true

- ➊ The system $\begin{pmatrix} A^\top & 0 \\ b^\top & 1 \end{pmatrix} \begin{pmatrix} y \\ \delta \end{pmatrix} = \begin{pmatrix} c \\ v \end{pmatrix}, \begin{pmatrix} y \\ \delta \end{pmatrix} \succeq 0$ has a solution.
 - Here, $y \in \mathbb{R}$ and $\delta \in \mathbb{R}$.
 - Implies dual is feasible and $OPT(dual) \leq v$

Formal Proof of Strong Duality

Primal LP

maximize $\langle c, x \rangle$
subject to $Ax \preceq b$

Dual LP

minimize $\langle b, y \rangle$
subject to $A^T y = c$
 $y \succeq 0$

Given $v \in \mathbb{R}$, by Farkas' Lemma exactly one of the following is true

① The system $\begin{pmatrix} A^T & 0 \\ b^T & 1 \end{pmatrix} \begin{pmatrix} y \\ \delta \end{pmatrix} = \begin{pmatrix} c \\ v \end{pmatrix}, \begin{pmatrix} y \\ \delta \end{pmatrix} \succeq 0$ has a solution.

- Here, $y \in \mathbb{R}$ and $\delta \in \mathbb{R}$.
- Implies dual is feasible and $OPT(dual) \leq v$

② The system $\begin{pmatrix} A & b \\ 0 & 1 \end{pmatrix} z \preceq 0, z^T \begin{pmatrix} c \\ v \end{pmatrix} > 0$ has a solution.

Let $z = \begin{pmatrix} z_1 \\ z_2 \end{pmatrix}$ be a solution, where $z_1 \in \mathbb{R}^n$ and $z_2 \in \mathbb{R}$

- Ⓐ If $z_2 \neq 0$, then $x = -z_1/z_2$ is primal feasible and $c^T x > v$. Therefore $OPT(primal) > v$.

Formal Proof of Strong Duality

Primal LP

maximize $\langle c, x \rangle$
subject to $Ax \preceq b$

Dual LP

minimize $\langle b, y \rangle$
subject to $A^T y = c$
 $y \succeq 0$

Given $v \in \mathbb{R}$, by Farkas' Lemma exactly one of the following is true

1 The system $\begin{pmatrix} A^T & 0 \\ b^T & 1 \end{pmatrix} \begin{pmatrix} y \\ \delta \end{pmatrix} = \begin{pmatrix} c \\ v \end{pmatrix}$, $\begin{pmatrix} y \\ \delta \end{pmatrix} \succeq 0$ has a solution.

- Here, $y \in \mathbb{R}$ and $\delta \in \mathbb{R}$.
- Implies dual is feasible and $OPT(dual) \leq v$

2 The system $\begin{pmatrix} A & b \\ 0 & 1 \end{pmatrix} z \preceq 0$, $z^T \begin{pmatrix} c \\ v \end{pmatrix} > 0$ has a solution.

Let $z = \begin{pmatrix} z_1 \\ z_2 \end{pmatrix}$ be a solution, where $z_1 \in \mathbb{R}^n$ and $z_2 \in \mathbb{R}$


- A If $z_2 \neq 0$, then $x = -z_1/z_2$ is primal feasible and $c^T x > v$. Therefore $OPT(primal) > v$.
- B If $z_2 = 0$, primal infeasible or unbounded, & dual infeasible (prove it)

Outline


- 1 Linear Programming Basics
- 2 Duality and Its Interpretations
- 3 Properties of Duals
- 4 Weak and Strong Duality
- 5 Consequences of Duality
- 6 Formal Proof of Strong Duality of LP
- 7 More Examples of Duality**

Shortest Path

Given a directed network $G = (V, E)$ where edge e has length $\ell_e \in \mathbb{R}_+$, find the minimum cost path from s to t .



Shortest Path



Primal LP


$$\begin{aligned} \min \quad & \sum_{e \in E} \ell_e x_e \\ \text{s.t.} \quad & \sum_{e \rightarrow v} x_e - \sum_{v \rightarrow e} x_e = \delta_v, \quad \forall v \in V. \\ & x_e \geq 0, \quad \forall e \in E. \end{aligned}$$

Dual LP

$$\begin{aligned} \max \quad & y_t - y_s \\ \text{s.t.} \quad & y_v - y_u \leq \ell_e, \quad \forall (u, v) \in E. \end{aligned}$$

Where $\delta_v = -1$ if $v = s$, 1 if $v = t$, and 0 otherwise.

Shortest Path



Primal LP

$$\begin{aligned} \min \quad & \sum_{e \in E} \ell_e x_e \\ \text{s.t.} \quad & \sum_{e \rightarrow v} x_e - \sum_{v \rightarrow e} x_e = \delta_v, \quad \forall v \in V. \\ & x_e \geq 0, \quad \forall e \in E. \end{aligned}$$

Dual LP

$$\begin{aligned} \max \quad & y_t - y_s \\ \text{s.t.} \quad & y_v - y_u \leq \ell_e, \quad \forall (u, v) \in E. \end{aligned}$$

Where $\delta_v = -1$ if $v = s$, 1 if $v = t$, and 0 otherwise.

Interpretation of Dual

Stretch s and t as far apart as possible, subject to edge lengths.

Maximum Weighted Bipartite Matching

Set B of buyers, and set G of goods. Buyer i has value w_{ij} for good j , and interested in at most one good. Find maximum value assignment of goods to buyers.

Maximum Weighted Bipartite Matching

Primal LP

$$\begin{array}{ll}\max & \sum_{i,j} w_{ij} x_{ij} \\ \text{s.t.} & \sum_{j \in G} x_{ij} \leq 1, \quad \forall i \in B. \\ & \sum_{i \in B} x_{ij} \leq 1, \quad \forall j \in G. \\ & x_{ij} \geq 0, \quad \forall i, j.\end{array}$$

Dual LP

$$\begin{array}{ll}\min & \sum_{i \in B} u_i + \sum_{j \in G} p_j \\ \text{s.t.} & u_i + p_j \geq w_{ij}, \quad \forall i \in B, j \in G. \\ & u_i \geq 0, \quad \forall i \in B. \\ & p_j \geq 0, \quad \forall j \in G.\end{array}$$

Maximum Weighted Bipartite Matching

Primal LP

$$\begin{array}{ll}\max & \sum_{i,j} w_{ij} x_{ij} \\ \text{s.t.} & \sum_{j \in G} x_{ij} \leq 1, \quad \forall i \in B. \\ & \sum_{i \in B} x_{ij} \leq 1, \quad \forall j \in G. \\ & x_{ij} \geq 0, \quad \forall i, j.\end{array}$$

Dual LP

$$\begin{array}{ll}\min & \sum_{i \in B} u_i + \sum_{j \in G} p_j \\ \text{s.t.} & u_i + p_j \geq w_{ij}, \quad \forall i \in B, j \in G. \\ & u_i \geq 0, \quad \forall i \in B. \\ & p_j \geq 0, \quad \forall j \in G.\end{array}$$

Interpretation of Dual

- p_j is price of good j
- u_i is utility of buyer i
- Complementary Slackness:
 - A buyer i only grabs goods j maximizing $w_{ij} - p_j$
 - Only fully assigned goods have non-zero price
 - A buyer with nonzero utility must receive an item

2-Player Zero-Sum Games

Rock-Paper-Scissors

	R	P	S
R	0	1	-1
P	-1	0	1
S	1	-1	0

- Two players, row and column
- Game described by matrix A
- When row player plays pure strategy i and column player plays pure strategy j , row player pays column player A_{ij}

2-Player Zero-Sum Games

Rock-Paper-Scissors

	R	P	S
R	0	1	-1
P	-1	0	1
S	1	-1	0

- Two players, row and column
- Game described by matrix A
- When row player plays pure strategy i and column player plays pure strategy j , row player pays column player A_{ij}
- **Mixed Strategy**: distribution over pure strategies

2-Player Zero-Sum Games

Rock-Paper-Scissors

	R	P	S
R	0	1	-1
P	-1	0	1
S	1	-1	0

- Two players, row and column
- Game described by matrix A
- When row player plays pure strategy i and column player plays pure strategy j , row player pays column player A_{ij}
- **Mixed Strategy**: distribution over pure strategies
- If one of the players moves first, the other observes his mixed strategy but not the outcome of his coin flips.

2-Player Zero-Sum Games

- Assume row player moves first with distribution $y \in \Delta_m$
 - Payment as a function of Column's strategy given by $y^\top A$
 - A best response by column is pure strategy j maximizing $(y^\top A)_j$
 - Row player solves an LP to determine optimal strategy y , payment u for himself

	x_1	x_2	x_3	x_4
y_1	a_{11}	a_{12}	a_{13}	a_{14}
y_2	a_{21}	a_{22}	a_{23}	a_{24}
y_3	a_{31}	a_{32}	a_{33}	a_{34}

2-Player Zero-Sum Games

- Assume row player moves first with distribution $y \in \Delta_m$
 - Payment as a function of Column's strategy given by $y^T A$
 - A best response by column is pure strategy j maximizing $(y^T A)_j$
 - Row player solves an LP to determine optimal strategy y , payment u for himself

Row Moves First

$$\begin{array}{ll}\min & \max_j \sum_i a_{ij} y_i \\ \text{s.t.} & \sum_i y_i = 1 \\ & y \succeq \vec{0}\end{array}$$

2-Player Zero-Sum Games

- Assume row player moves first with distribution $y \in \Delta_m$
 - Payment as a function of Column's strategy given by $y^T A$
 - A best response by column is pure strategy j maximizing $(y^T A)_j$
 - Row player solves an LP to determine optimal strategy y , payment u for himself

Row Moves First

min u

s.t.

$$\sum_i a_{ij} y_i \leq u, \quad \forall j.$$

$$\sum_i y_i = 1$$

$$y \succeq \vec{0}$$

2-Player Zero-Sum Games

- Assume row player moves first with distribution $y \in \Delta_m$
 - Payment as a function of Column's strategy given by $y^T A$
 - A best response by column is pure strategy j maximizing $(y^T A)_j$
 - Row player solves an LP to determine optimal strategy y , payment u for himself
- Similarly when column moves first, column solves an LP to determine optimal strategy x , payment v for himself

Row Moves First

$$\min \quad u$$

s.t.

$$\sum_i a_{ij} y_i \leq u, \quad \forall j.$$

$$\sum_i y_i = 1$$

$$y \succeq \vec{0}$$

Column Moves First

$$\max \quad v$$

s.t.

$$\sum_j a_{ij} x_j \geq v, \quad \forall i.$$

$$\sum_j x_j = 1$$

$$x \succeq \vec{0}$$

2-Player Zero-Sum Games

- Assume row player moves first with distribution $y \in \Delta_m$
 - Payment as a function of Column's strategy given by $y^T A$
 - A best response by column is pure strategy j maximizing $(y^T A)_j$
 - Row player solves an LP to determine optimal strategy y , payment u for himself
- Similarly when column moves first, column solves an LP to determine optimal strategy x , payment v for himself

Row Moves First

$$\min \quad u$$

s.t.

$$\sum_i a_{ij} y_i \leq u, \quad \forall j.$$

$$\sum_i y_i = 1$$

$$y \succeq \vec{0}$$

Column Moves First

$$\max \quad v$$

s.t.

$$\sum_j a_{ij} x_j \geq v, \quad \forall i.$$

$$\sum_j x_j = 1$$

$$x \succeq \vec{0}$$

These two optimization problems are LP Duals!

Duality and Zero Sum Games

Weak Duality

- $u \geq v$
- Zero sum games have a second mover advantage

Duality and Zero Sum Games

Weak Duality

- $u \geq v$
- Zero sum games have a second mover advantage

Strong Duality (Minimax Theorem)

- $u^* = v^*$
- There is no second or first mover advantage in zero sum games with mixed strategies
- Each player can guarantee $u^* = v^*$ even if they move first (i.e., regardless of other's strategy).
- y^*, x^* are simultaneously best responses to each other (Nash Equilibrium)

Duality and Zero Sum Games

Weak Duality

- $u \geq v$
- Zero sum games have a second mover advantage

Strong Duality (Minimax Theorem)

- $u^* = v^*$
- There is no second or first mover advantage in zero sum games with mixed strategies
- Each player can guarantee $u^* = v^*$ even if they move first (i.e., regardless of other's strategy).
- y^*, x^* are simultaneously best responses to each other (Nash Equilibrium)

Complementary Slackness

x^* randomizes over pure best responses to y^* , and vice versa.


Saddle Point Interpretation

Consider the matching pennies game

	H	T
H	-1	1
T	1	-1

- Unique equilibrium: each player randomizes uniformly
- If row player deviates, he pays out (weakly) more
- If column player deviates, he gets paid (weakly) less

Saddle Point Interpretation



- Unique equilibrium: each player randomizes uniformly
- If row player deviates, he pays out (weakly) more
- If column player deviates, he gets paid (weakly) less